

Part IV Overview. National implications and a science strategy for the way forward

Authors: S. Jeffress Williams, USGS; E. Robert Thieler, USGS; Benjamin T. Gutierrez, USGS; Dean B. Gesch, USGS; Donald R. Cahoon, USGS; K. Eric Anderson, USGS

Climate change and effects such as sea-level rise have global implications and will increasingly affect the entire nation. While this Product focuses primarily on the mid-Atlantic region of the United States, many of the issues discussed in earlier chapters are relevant at the national scale.

Chapter 13 draws on findings from the mid-Atlantic focus area that have relevance to other parts of the United States, provides an overview of coastal environments and landforms in the United States, and describes the issues faced in understanding how these environments may be impacted and respond to sea-level rise. The diversity of U.S. coastal settings includes bedrock coasts in Maine, glacial bluffs in New York, barrier islands in the mid-Atlantic and Gulf of Mexico, coral reefs in Florida, the Caribbean, and Hawaii, one of the world's major delta systems in Louisiana, a wide variety of pocket beaches and cliffed coasts along the Pacific coast, Pacific atolls, and a number of arctic coastline types in Alaska. In addition, the large bays and estuaries around the country also exhibit a diverse range of shoreline types, large wetland systems, and extensive coastal habitats.

8185

8186 Understanding how the different coastal environments of the United States will respond
8187 to future climate and sea-level change is a major challenge. In addition, as highlighted in
8188 earlier Parts of this Product, human actions and policy decisions also substantially
8189 influence the evolution of the coast. The knowledge gaps and data limitations identified
8190 in this Product focusing on the mid-Atlantic have broad relevance to the rest of the U.S.
8191 Chapter 14 identifies opportunities for increasing the scientific understanding of future
8192 sea-level rise impacts. This includes basic and applied research in the natural and the
8193 social sciences. A significant emphasis is placed on developing linkages between
8194 scientists, policy makers, and stakeholders at all levels, so that information can be shared
8195 and utilized efficiently and effectively as sea-level rise mitigation and adaptation plans
8196 evolve.

8197

Chapter 13. Implications of Sea-Level Rise to the Nation

Authors: S. Jeffress Williams, USGS; Benjamin T. Gutierrez, USGS; James G. Titus, USEPA; Eric Anderson, USGS (retired); Stephen K. Gill, NOAA; Donald R. Cahoon, USGS; E. Robert Thieler, USGS; and Dean B. Gesch, USGS

KEY FINDINGS

- Nationwide, more than one-third of the United States population currently lives in the coastal zone and movement to the coast and development continues, along with the current and growing vulnerability to coastal hazards such as storms and sea-level rise. Fourteen of the 20 largest U.S. urban centers are located along the coast. With the very likely accelerated rise in sea level and increased storm intensity, the conflicts between people and development at the coast and the natural processes will increase, causing economic and societal impacts.
- For much of the U.S., shores comprised of barrier islands, dunes, spits, and sandy bluffs, erosion processes will dominate at highly variable rates in response to sea-level rise and storms over the next century and beyond. Some coastal landforms in the U.S. may undergo large changes in shape and location if the rate of sea-level rise increases as predicted. Increased inundation and more frequent flooding will affect estuaries and low-lying coastal areas. The response to these driving forces will vary depending on the type of coastal landform and local conditions, but will

- 8220 be more extreme, more variable and less predictable than the changes observed
8221 over the last century.
- 8222 • For higher sea-level rise scenarios, some barrier island coasts and wetlands may
8223 cross thresholds and undergo significant and irreversible changes. These changes
8224 include rapid landward migration and segmentation of some barrier islands and
8225 disintegration and drowning of wetlands.
 - 8226 • Nationally, tidal wetlands already experiencing submergence by sea-level rise and
8227 associated land loss, in concert with other factors, will continue to deteriorate in
8228 response to changing climate.
 - 8229 • Coastal change is driven by complex and interrelated processes. Over the next
8230 century and beyond, with an expected acceleration in sea-level rise, the potential
8231 for coastal change is likely to be greater than has been observed in historic past.
8232 These changes to coastal regions will have especially large impacts on urban
8233 centers and developed areas. Some portions of the U.S. coast, however, will be
8234 subject primarily to inundation from sea-level rise over the next century. A
8235 substantial challenge remains to quantify the various effects of sea-level rise and
8236 to identify the dominant coastal change processes for each region of the U.S.
8237 coast.
 - 8238 • Many coastal areas in the United States will likely experience an increased
8239 frequency and magnitude of storm-surge flooding and coastal erosion due to
8240 storms over the next century, in response to sea-level rise. The impacts from these
8241 storm events are likely to extend farther inland from the coast than those that
8242 would be affected by sea-level rise alone.

8243 • Understanding, predicting, and responding to the environmental and societal
8244 effects of sea-level rise would benefit from a national program of integrated
8245 research that includes the natural and social sciences. Research on adaptation,
8246 mitigation, and avoidance-of-risk measures would enable improved understanding
8247 of the many and varied potential societal impacts of sea-level rise that would
8248 benefit the U.S as well as coastal nations around the world.

8249

8250 **13.1 INTRODUCTION**

8251 As defined in the SAP 4.1 Prospectus and discussed in earlier chapters, this Product
8252 focuses on assessing potential impacts to the mid-Atlantic region; however, some
8253 discussion of impacts to other regions and the nation as a whole is warranted. The mid-
8254 Atlantic region is highly vulnerable to sea-level rise, but regions like the central Gulf
8255 Coast (Louisiana, Texas) are just as vulnerable or more so. The challenge in carrying out
8256 a national assessment is that nationwide data bases and scientific publications of national
8257 scale and scope are limited. Modest efforts at monitoring and observations for national-
8258 scale assessments of coastal change and hazards are underway by various organizations,
8259 but more effort is needed. The discussion in ection 13.3 is largely the expert opinions of
8260 the lead authors, informed by results of the two expert science panel reports (Reed *et al.*,
8261 2008, Gutierrez, *et al.*, 2007) and available scientific literature. Because of the relative
8262 lack of adequate background literature and high reliance on expert opinion, the likelihood
8263 statements as used in other chapters are not included in this discussion of potential
8264 impacts to the nation.

8265

A large and expanding proportion of the U.S. population and related urban development is located along the Atlantic, Gulf of Mexico, and Pacific coasts and increasingly conflicts with the natural processes associated with coastal change from storms and sea-level rise (see a review in Williams *et al.*, 1991). Development in low-lying regions (*i.e.*, New Orleans) and islands (*e.g.*, in the Chesapeake Bay, Caribbean, Pacific Ocean) are particularly at risk (see Gibbons and Nicholls, 2006). In the future, as the effects of climate change intensify, these interactions will become more frequent and more challenging to society. Currently, more than one-third of the population lives in the coastal zone and movement to the coast and development continues, along with the growing vulnerability to coastal hazards. Fourteen of the 20 largest U.S. urban centers are located along the coast (Crossett *et al.*, 2004; Crowell *et al.*, 2007). With the likely accelerated rise in sea level and increased storm intensity, the conflicts between people and development at the coast and the natural processes will increase, affecting all parts of society (Leatherman, 2001; FitzGerald *et al.*, 2008).

Global sea-level rise associated with climate change is likely to be in the range of 19 cm to as much as 1 m over the next century and possibly as much as 4 to 6 m over the next several centuries (IPCC, 2007; Rahmstorf, 2007; Rahmstorf, *et al.*, 2007; Overpeck *et al.*, 2006). The expected rise will increase erosion and the frequency of flooding and coastal areas will be at increasing risk. For some regions, adaptation using engineering means may be effective; for other coastal areas, however, adaption by relocation landward to higher elevated ground may be appropriate for longer-term sustainability (NRC, 1987).

Coastal landforms reflect the complex interaction between the natural physical processes that act on the coast, the geologic characteristics of the coast, and human activities. Spatial and temporal variations in these physical processes and the geology along the coast are responsible for the wide variety of landforms around the United States (Williams, 2003). With future sea-level rise, portions of the U.S. ocean coast are likely to undergo long-term net erosion, at rates higher than those that have been observed over the past century (see Chapter 3). The exact manner and rates at which these changes are likely to occur depend on the character of coastal landforms (*e.g.*, barrier islands, cliffs) and the physical processes (*e.g.*, waves and winds) that shape these landforms (see Chapters 3 and 4). Low-relief coastal regions, areas undergoing land subsidence and land subject to frequent storm landfalls, such as the northern Gulf of Mexico, Florida, Hawaii, Puerto Rico, the San Francisco-Sacramento Delta region, and the Mid-Atlantic region, are particularly vulnerable.

13.2 TYPES OF COASTS

Coasts are dynamic junctions of the oceans, atmosphere, and land and differ greatly in physical character and vulnerability to erosion, storms, and sea-level rise (NRC, 1990). The principal coastal types are described in Chapters 3 and 4, and summarized below. With future sea-level rise, all of these landforms will become more dynamic (Nicholls *et al.*, 2007), but predicting and quantifying changes that are likely to occur with high confidence is currently scientifically challenging.

13.2.1 Cliff and Bluff Shorelines

Substantial portions of the U.S. coast are comprised of coastal cliffs and bluffs that vary greatly in height, morphology, and sedimentary composition. These occur predominantly along the New England and Pacific coasts, Hawaii, and Alaska. Coastal cliff is a general term that refers to steep slopes along the shoreline that commonly form in response to long-term rise in sea-level. The term “bluff” also can refer to escarpments eroded into unlithified material, such as glacial till, along the shore (Hampton and Griggs, 2004). The terms “cliff” and “bluff” are often used interchangeably. Coastal cliffs erode in response to a variety of both marine and terrestrial processes. Cliff retreat can be fairly constant, but can also be episodic. In contrast to sandy coasts, which may erode landward or accrete seaward, cliffs retreat only in a landward direction. Because rocky cliff coasts are composed of resistant materials, erosion can occur more slowly than for those comprised of unconsolidated sediments and response times to sea-level rise much longer than for sandy coasts (NRC, 1987), but land slumping due to wave action or land surface water runoff can result in rapid retreat. Hampton and Griggs (2004) provide a review of the origin, U.S. distribution, evolution, and regional issues associated with coastal cliffs. Predicting the response of coastal cliffs to future sea-level rise is a topic of active research (Trenhaile, 2001; Walkden and Hall, 2005; Dickson *et al.*, 2007; Walkden and Dickson, 2008)

13.2.2 Sandy Shores, Pocket Beaches, Barrier Beaches, Spits, and Dunes

Sandy beaches are often categorized into a few basic types which commonly include mainland, pocket, and barrier beaches (Wells, 1995; Davis and FitzGerald, 2004). The sediments that comprise beaches are derived mainly from the erosion of the adjacent

mainland and continental shelf, and sometimes from sediments supplied from coastal rivers. Mainland beaches occur where the land intersects the shore. Some mainland beaches occur in low-relief settings and are surrounded by coastal dunes, while others occur along steep portions of the coast and are backed by bluffs. Examples of mainland beaches include the shores of eastern Long Island, northern New Jersey (Oertel and Kraft, 1994), and parts of Delaware, (Kraft, 1971). Pocket beaches form in small bays, often occurring between rocky headlands and are common along parts of the southern New England coast, portions of California and Oregon (Hapke *et al.*, 2006), and in parts of the Hawaiian Islands. Barrier beaches and spits are the most abundant coastal landforms along the Atlantic and Gulf of Mexico coasts. In general, it is expected that accelerations in sea-level rise will enhance beach erosion globally, but on a local scale this response will depend on the sediment budget (Nicholls *et al.*, 2007).

13.2.3 Coastal Marshes, Mangroves, and Mud Flat Shorelines

Coastal wetlands include swamps and tidal flats, salt and brackish marshes, mangroves, and bayous. They form in low-relief, low-energy sheltered coastal environments, often in conjunction with river deltas, landward of barrier islands, and along the flanks of estuaries (*e.g.*, Delaware Bay, Chesapeake Bay, Everglades, Lake Pontchartrain, Galveston Bay, San Francisco Bay, and Puget Sound). Most coastal wetlands are in Louisiana, North and South Carolina, south Florida, and Alaska (Dahl, 1990; NRC, 1995a). Wetlands are extremely vulnerable to sea-level rise and can maintain their elevation and viability only if sediment accumulation (both mineral and organic matter) keeps pace with sea-level rise (Cahoon *et al.*, 2006; Nyman *et al.*, 2006; Morris *et al.*,

2002; Rybczyk and Cahoon, 2002). Future wetland area will also be determined, in part, by the amount of space (*e.g.*, mud flat or tidal flat area) available for landward migration and the rates of lateral erosion of the seaward edge of the marsh (see Chapter 4; Poulter, 2005). Wetlands will be especially vulnerable to the higher projected rates of future sea-level rise (*e.g.*, greater than 70 cm by the year 2100), but some will survive a 1 meter rise (Morris *et al.*, 2002). Even under lower accelerated sea-level rise rates, wetlands may be sustained only where conditions are optimal for vertical wetland development (*e.g.*, abundant sediment supply and low regional subsidence rate) (Rybczyk and Cahoon, 2002).

Mud flat shorelines represent a relatively small portion of U.S. coasts, but are important in providing the foundation for wetlands and marshes (Mitsch and Gosselink, 1986). They are frequently associated with wetlands, and occur predominately in low-energy, low-relief regions with high inputs of fine-grained river-born sediments and organic materials and large tidal ranges. These shoreline types are common in western Louisiana (*i.e.*, Chenier Plain) and along northeastern parts of the Gulf Coast of Florida. Muddy coasts may be drowned with sea-level rise unless sediment inputs are sufficiently large, such as the Atchafalaya River delta region of southwestern Louisiana, and the flats are able to be colonized by plants.

13.2.4 Tropical Coral Reef Coasts

Tropical coral reefs, made up of living organisms very sensitive to ocean temperature and chemistry, are found in the U.S. along the south coast of Florida; around the Hawaiian

Islands, Puerto Rico, the Virgin Islands, and many of the U.S. territories in the Pacific (Riegl and Dodge, 2008). In tropical environments, living coral organisms build reefs that are important ecological resources (Smith and Buddemeier, 1992; Boesch *et al.*, 2000). Most corals are able to tolerate rates of sea-level rise of 10 to 20 mm per year or more (Smith and Buddemeier, 1992; Bird, 1995; Wells, 1995; Hallock, 2005). Nonetheless, the ability of coral reef systems to survive future sea-level rise will depend heavily on other climate change impacts such as increase in ocean temperature and or acidity, sediment runoff from the land, as well as episodic storm erosion (Hallock, 2005; Nicholls *et al.*, 2007). In addition, human caused stresses such as over-fishing or pollution can contribute to the vulnerability of these systems to climate change (Buddemeier *et al.*, 2004; Mimura *et al.*, 2007).

13.3 Potential for Future Shoreline Change

Over the next century and beyond, with an expected acceleration in sea-level rise, the potential for coastal change will increase and coastal change is likely to be more widespread and variable than has been observed in historic past (NRC, 1987; Brown and McLachlan, 2002; Nicholls *et al.*, 2007). However, it is difficult at present to quantitatively attribute shoreline changes directly to sea-level rise (Rosenzweig *et al.*, 2007). The potential changes include increased coastal erosion, more frequent tidal and storm-surge flooding of low-relief areas, and wetland deterioration and losses. Many of these changes will occur in all coastal states. These changes to the coastal zone can be expected to have especially large impacts to developed areas (Nicholls *et al.*, 2007). Some portions of the U.S. coast will be subject principally to inundation from sea-level

rise over the next century, including upper reaches of bays and estuaries (*e.g.*, Chesapeake and Delaware Bays, Tampa Bay, Lake Pontchartrain, San Francisco Bay), and hardened urban shorelines. Erosion, sediment transport, and sediment deposition in coastal environments are active processes and will drive coastal change in concert with the combined effects of future sea-level rise and storms (Stive, 2004).

Coastal landforms may become even more dynamic and that erosion will dominate changes in shoreline position over the next century and beyond (Nicholls *et al.*, 2007). Wetlands with sufficient sediment supply and available land for inland migration may be able to maintain elevation, keeping pace with sea-level rise, but sediment starved wetlands and those constrained by engineering structures (*e.g.*, seawalls, revetments) or steep uplands are likely to deteriorate and convert to open water through vertical accretion deficits and lateral erosion (see Chapter 4). On barrier island shores, erosion is likely to occur on both the ocean front and the landward side of the island due to a combination of storm activity, changes in sediment budget, more frequent tidal flooding, and rising water levels (Nicholls *et al.*, 2007).

Sea-level rise is a particular concern for islands (Mimura *et al.*, 2007). Of particular concern are islands comprised of coral atolls (*e.g.*, Midway Atoll), which are typically low-lying and dependent on the health of coral reefs that fringe the atolls. Populated islands with higher elevations (*e.g.*, the Northern Mariana Islands) are also frequently at risk as the infrastructure is frequently located in low-lying coastal regions along the periphery of the islands.

8427

8428 Many coastal areas in the United States will likely experience an increased frequency and
8429 magnitude of storm-surge flooding, greater wave heights, and more erosion due to storms
8430 as part of the response to sea-level rise (NRC, 1987; Woodworth and Blackman, 2004;
8431 Nicholls *et al.*, 2007; Gutowski *et al.*, 2008). Impacts from these storm events may
8432 extend farther inland than those that would be affected by sea-level rise alone. Many
8433 regions may also experience large changes to coastal systems, such as increased rates of
8434 erosion, barrier island and dune landward migration, and potential barrier island collapse
8435 (Nicholls *et al.*, 2007; see also Chapters 1, 3, and 14 for discussion of geomorphic
8436 thresholds). The potential of crossing thresholds, potentially leading to barrier and
8437 wetland collapse, may increase with higher rates of sea-level rise.

8438

8439 The use of so called “soft” coastal engineering mitigation measures, such as beach
8440 nourishment, usually using sand dredged from offshore Holocene-age sand bodies, may
8441 reduce the risk of storm flooding and coastal erosion temporarily (NRC, 1987, 1995b).
8442 However, an important issue is whether or not these practices are able to be maintained
8443 into the future to provide sustainable and economical shoreline protection in the face of
8444 high cost, need for periodic re-nourishment, and limited sand resources of suitable quality
8445 for nourishment for many regions of the country (NRC, 1995b; Magoon *et al.*, 2004).
8446 Results from offshore geologic mapping studies indicate that most continental shelf
8447 regions of the U.S. have relatively limited Holocene-age sediment that can be deemed
8448 available and suitable for uses such as beach nourishment (Schwab *et al.*, 2000; Gayes *et*
8449 *al.*, 2003; Pilkey *et al.*, 1981; Kraft, 1971). In some cases, potential sand volumes are

reduced because of economic and environmental factors such as water depth, benthic environmental concerns, and concerns that sand removal may alter sediment exchange with the adjacent coast (Bliss *et al.*, 2009). The result is limited volumes of high-quality offshore sand resources readily available for beach nourishment. The issue of relying long term on using offshore sand for beach nourishment to mitigate erosion is important and needs to be addressed.

More widespread implementation of regional sediment or best sediment management practices to conserve valuable coastal sediments from offshore disposal of clean sandy dredged spoils will enhance the long term sustainability of sandy coastal landforms (NRC, 2007). The use of so called “hard” engineering structures (*e.g.*, seawalls, breakwaters) to protect property from erosion and flooding may be justified for urban coasts, but their use on sandy shores can further exacerbate erosion over time due to disruption of sediment transport processes. Alternatives, such as relocation landward, strategic removal of development or limiting redevelopment following storm disasters from highly vulnerable parts of the coast, may provide longer term sustainability of both coastal landforms and development, especially if the higher rates of sea-level rise are realized (NRC, 1987). An example of abandonment of an island in Chesapeake Bay due to sea-level rise is detailed in Gibbons and Nicholls (2006). If coastal development is relocated, those areas could be converted to marine protected areas, public open-space lands that would serve to buffer sea-level rise effects landward and also provide recreation benefits and wildlife habitat values (see Salm and Clark, 2000).

13.4 CONCLUSIONS

Global climate is changing, largely due to carbon emissions from human activities (IPCC, 2001; 2007). Sea-level rise is one of the impacts of climate change that will affect all coastal regions of the United States over the next century and beyond (NRC, 1987; Nicholls *et al.*, 2007). The scientific tools and techniques for assessing the effects of future sea-level rise on coastal systems are improving, but much remains to be done in order to develop useful forecasts of potential effects. Chapter 14 of this Product identifies research opportunities that, if implemented, would lead to better understanding and prediction of sea-level rise effects that are likely to further impact the United States in the near future. Planning for accelerating sea-level rise should include thorough evaluation of a number of alternatives, such as cost-effective and sustainable shore protection and strategic relocation of development within urban centers. Important decisions like these should ideally be based on the best available science and careful consideration of long-term benefits for a sustainable future, and the total economic, social, and environmental costs of various methods of shore protection, relocation, and adaptation.

8489 CHAPTER 13 REFERENCES

- 8490 **Bird**, E.C.F., 1995: Present and future sea-level: the effects of predicted global changes.
8491 In: *Climate Change: Impact and Coastal Habitation*, [Eisma, D. (ed.)]. CRC
8492 Press, Boca Raton, FL, pp. 29-56.
- 8493 **Bliss**, J.D., S.J. Williams, and K.S. Bolm, 2009: Modeling and assessment of marine sand
8494 resources, New York Bight, USA. In: *Contributions to industrial minerals*
8495 *research*. [Bliss, J.D., P.R. Moyle, and K.R. Long, (eds.)]. U.S. Geological
8496 Survey bulletin 2209-M. U.S. Geological Survey, [Reston, VA], 22 pp.
- 8497 **Boesch**, D.F., J.C. Field, and D. Scavia (eds.), 2000: *The Potential Consequences of*
8498 *Climate Variability and Change on Coastal Areas and Marine Resources*. NOAA
8499 Coastal Ocean Program decision analysis series #21. National Oceanic and
8500 Atmospheric Administration, Silver Spring, MD, 163 pp.
8501 <<http://www.cop.noaa.gov/pubs/das/das21.pdf>>
- 8502 **Brown**, A.C. and A. McLachlan, 2002: Sandy shore ecosystems and the threats facing
8503 them: some predictions for the year 2025. *Environmental Conservation*, **29(1)**,
8504 62-77.
- 8505 **Buddemeier**, R.W., J.A. Kleypas, and R.B. Aronson, 2004: *Coral Reefs and Goba*
8506 *Climate Change*. Pew Center on Global Climate Change, Arlington, VA, 44 pp.
- 8507 **Cahoon**, D.R., P.F. Hensel, T. Spencer, D.J. Reed, K.L. McKee, and N. Saintilan, 2006:
8508 Coastal wetland vulnerability to relative sea level rise: wetland elevation trends
8509 and process controls. In: *Wetlands and Natural Resource Management*
8510 [Verhoeven, J.T.A., B. Beltman, R. Bobbink, and D. Whigham (eds.)]. Ecological
8511 studies volume 190, Springer, Berlin and New York, pp. 271-292.
- 8512 **Crossett**, K., T.J. Culliton, P. Wiley, and T.R. Goodspeed, 2004: *Population Trends*
8513 *along the Coastal United States, 1980–2008*. National Oceanic and Atmospheric
8514 Administration, Silver Spring, MD, 47 pp.

- 8515 **Crowell**, M., K. Coulton, and S. McAfee, 2007: How many people live in coastal areas?.
- 8516 *Journal of Coastal Research*, **23(5)**, iii-vi, editorial.
- 8517 **Dahl**, T.E., 1990: *Wetlands losses in the United States 1780's to 1980's*, U.S. Department
- 8518 of the Interior, Fish and Wildlife Service, Washington, DC, 21 pp.
- 8519 **Davis**, R.A. and D.M. FitzGerald, 2004: *Beaches and Coasts*. Blackwell Publishing,
- 8520 Malden, MA, 419 pp.
- 8521 **Dickson**, M.E., M.J.A. Walkden, and J.W. Hall, 2007: Systemic impacts of climate
- 8522 change on an eroding coastal region over the twenty-first century. *Climatic*
- 8523 *Change*, **84(2)**, 141-166.
- 8524 **FitzGerald**, D.M., M.S. Fenster, B.A. Argow, and I.V. Buynevich, 2008: Coastal
- 8525 impacts due to sea-level rise. *Annual Reviews of Earth and Planetary Sciences*,
- 8526 **36**, 601-647.
- 8527 **Gayes**, P.T., W.C. Schwab, N.W. Driscoll, R.A. Morton, W.E. Baldwin, J.F. Denny, E.E.
- 8528 Wright, M.S. Harris, M.P. Katuna, T.R. Putney, and E. Johnstone, 2003:
- 8529 Sediment dispersal pathways and conceptual sediment budget for a sediment
- 8530 starved embayment; Long Bay, South Carolina. In: *Coastal Sediments '03, 5th*
- 8531 *Annual Symposium on Coastal Engineering and Science of Coastal Sediment*
- 8532 *Processes*, Clearwater Beach, FL, May 18-23, 2003. East Meets West
- 8533 Productions, Corpus Christi, TX, 14 pp.
- 8534 **Gibbons**, S.J.A. and R.J. Nicholls, 2006: Island abandonment and sea-level rise: an
- 8535 historical analog from the Chesapeake Bay, USA. *Global Environmental Change*,
- 8536 **16(1)**, 40-47.
- 8537 **Gutierrez**, B.T., S.J. Williams, and E.R. Thieler, 2007: *Potential for Shoreline Changes*
- 8538 *Due to Sea-level Rise along the U.S. Mid-Atlantic Region*. U.S. Geological Survey
- 8539 open file report 2007-1278. U.S. Geological Survey, Reston, VA, 26 pp.
- 8540 <<http://pubs.usgs.gov/of/2007/1278/>>

- 8541 **Gutowski**, W.J., G.C. Hegerl, G.J. Holland, T.R. Knutson, L.O. Mearns, R.J. Stouffer,
8542 P.J. Webster, M.F. Wehner, and F.W. Zwiers, 2008: Causes of observed changes
8543 in extremes and projections of future changes. In: *Weather and Climate Extremes*
8544 *in a Changing Climate: Regions of Focus: North America, Hawaii, Caribbean,*
8545 *and U.S. Pacific Islands* [Karl, T.R., G.A. Meehl, C.D. Miller, S.J. Hassol, A.M.
8546 Waple, and W.L. Murray (eds.)]. Synthesis and Assessment Product 3.3. U.S.
8547 Climate Change Science Program, Washington, DC, pp. 81-116.
- 8548 **Hallock**, P., 2005: Global change and modern coral reefs: new opportunities to
8549 understand shallow-water carbonate depositional processes. *Sedimentary*
8550 *Geology*, **175**(1-4), 19-33.
- 8551 **Hampton**, M.A. and G.B. Griggs (eds.), 2004: *Formation, Evolution, and Stability of*
8552 *Coastal Cliffs — Status and Trends*. U.S. Geological Survey professional paper
8553 1693. U.S. Geological Survey, Reston, VA, 123 pp.
- 8554 **IPCC** (Intergovernmental Panel on Climate Change), 2001: *Climate Change 2001: The*
8555 *Scientific Basis*. Contribution of Working Group I to the Third Assessment Report
8556 of the Intergovernmental Panel on Climate Change [Houghton, J.T., Y. Ding, D.J.
8557 Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson
8558 (eds.)]. Cambridge University Press, Cambridge, UK, and New York, 881 pp.
- 8559 **IPCC** (Intergovernmental Panel on Climate Change), 2007: *Climate Change 2007: The*
8560 *Physical Science Basis*. Contribution of Working Group I to the Fourth
8561 Assessment Report of the Intergovernmental Panel on Climate Change [Solomon,
8562 S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, and H.L.
8563 Miller (eds.)]. Cambridge University Press, Cambridge, UK and New York, 996
8564 pp.
- 8565 **Leatherman**, S.P., 2001: Social and economic costs of sea level rise. In: *Sea Level Rise:*
8566 *History and Consequences* [Douglas, B.C., M.S. Kearney, and S.P. Leatherman
8567 (eds.)]. Academic Press, San Diego, pp. 181-223.

- 8568 **Magoon**, O.T., S.J. Williams, L.K. Lent, S.L. Douglass, B.L. Edge, J.A. Richmond, D.D.
8569 Treadwell, L.C. Ewing, and A.P. Pratt, 2004: Economic impacts of anthropogenic
8570 activities on coastlines of the U.S., In: *Coastal Engineering 2004*, Proceedings of
8571 the 29th International Conference, Lisbon, Portugal. World Scientific Publishing,
8572 New Jersey and London, pp. 3022-3035.
- 8573 **Mimura**, N.L., L. Nurse, R.F. McLean, J. Agard, L. Briguglio, P. Lefale, R. Payet, and
8574 G. Sem, 2007: Small islands. In: *Climate Change 2007: Impacts, Adaptation and*
8575 *Vulnerability*. Contribution of Working Group II to the Fourth Assessment Report
8576 of the Intergovernmental Panel on Climate Change [Parry, M.L., O.F. Canziani,
8577 J.P., Palutikof, P.J. van der Linden, and C.E. Hanson (eds.)]. Cambridge
8578 University Press, Cambridge, UK, and New York, pp. 688-716.
- 8579 **Mitsch**, W.J. and J.G. Gosselink, 1986: *Wetlands*. Van Nostrand, New York, 537 pp.
- 8580 **Morris**, J.T., P.V. Sundareshwar, C.T. Nietch, B. Kjerfve, and D.R. Cahoon, 2002:
8581 Responses of coastal wetlands to rising sea level. *Ecology*, **83(10)**, 2869-2877.
- 8582 **NRC** (National Research Council), 1987: *Responding to Changes in Sea Level*. National
8583 Academy Press, Washington, DC, 148 pp.
- 8584 **NRC** (National Research Council), 1990: *Managing Coastal Erosion*. National Academy
8585 Press, Washington, DC, 182 pp.
- 8586 **NRC** (National Research Council), 1995a: *Wetlands: Characteristics and Boundaries*.
8587 National Academy Press, Washington, DC, 306 pp.
- 8588 **NRC** (National Research Council), 1995b: *Beach Nourishment and Protection*. National
8589 Academy Press, Washington, DC, 334 pp.
- 8590 **NRC** (National Research Council), 2007: *Mitigating Shore Erosion Along Sheltered*
8591 *Coasts*. National Academies Press, Washington, DC, 174 pp.
- 8592 **Nicholls**, R.J., P.P. Wong, V.R. Burkett, J.O. Codignotto, J.E. Hay, R.F. McLean, S.
8593 Ragoonaden, and C.D. Woodroffe, 2007: Coastal systems and low-lying areas. In:

- 8594 *Climate Change 2007: Impacts, Adaptation and Vulnerability*. Contribution of
8595 Working Group II to the Fourth Assessment Report of the Intergovernmental
8596 Panel on Climate Change [Parry, M.L., O.F. Canziani, J.P. Palutikof, P.J. van der
8597 Linden, and C.E. Hanson (eds.)]. Cambridge University Press, Cambridge, UK,
8598 and New York, pp. 315-356.
- 8599 **Nyman**, J.A., R.J. Walters, R.D. Delaune, and W.H. Patrick Jr., 2006: Marsh vertical
8600 accretion via vegetative growth. *Estuarine Coastal and Shelf Science*, **69(3-4)**,
8601 370-380.
- 8602 **Oertel**, G.F. and J.C. Kraft, 1994: New Jersey and Delmarva barrier islands. In: *Geology*
8603 *of Holocene Barrier Island Systems* [Davis, R.A. (ed.)]. Springer-Verlag, New
8604 York, pp. 207-232.
- 8605 **Overpeck**, J.T., B.L. Otto-Bliesner, G.H. Miller, D.R. Muhs, R.B. Alley, and J.T. Keihl,
8606 2006: Paleo-climatic evidence for the future ice-sheet instability and rapid sea
8607 level rise. *Science*, **311(5768)**, 1747-1750.
- 8608 **Pilkey**, O.H., B.W. Blackwekder, H.J. Knebel, and M.W. Ayers, 1981: The Georgia
8609 Embayment continental shelf: Stratigraphy of a submergence. *Geological Society*
8610 *of America Bulletin*, **92(1)**, 52-63.
- 8611 **Rahmstorf**, S., 2007: A semi-empirical approach to projecting future sea-level rise.
8612 *Science*, **315(5810)**, 368-370.
- 8613 **Rahmstorf**, S., A. Cazenave, J.A. Church, J.E. Hansen, R.F. Keeling, D.E. Parker, and
8614 R.C.J. Somerville, 2007: Recent climate observations compared to projections.
8615 *Science*, **316(5825)**, 709.
- 8616 **Reed**, D.J., D. Bishara, D. Cahoon, J. Donnelly, M. Kearney, A. Kolker, L. Leonard,
8617 R.A. Orson, and J.C. Stevenson, 2008: Site-specific scenarios for wetlands
8618 accretion as sea level rises in the mid-Atlantic region. Section 2.1 in: *Background*
8619 *Documents Supporting Climate Change Science Program Synthesis and*
8620 *Assessment Product 4.1: Coastal Elevations and Sensitivity to Sea Level Rise*

- 8621 [Titus, J.G. and E.M. Strange (eds.)]. EPA 430R07004. U.S. Environmental
8622 Protection Agency, Washington, DC, pp. 134-174.
8623 <<http://epa.gov/climatechange/effects/coastal/background.html>>
- 8624 **Riegl**, B. and E. Dodge (eds.), 2008: *Coral Reefs of the USA*. Coral reefs of the
8625 worldvolume 1. Springer-Verlag, Dordrecht and London, 806 pp.
- 8626 **Rosenzweig**, C., G. Casassa, D.J. Karoly, A. Imeson, C. Liu, A. Menzel, S. Rawlins, T.L.
8627 Root, B. Seguin, P. Tryjanowski, and C.E. Hanson, 2007: Assessment of observed
8628 changes and responses in natural and managed systems. In: *Climate Change*
8629 *2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II*
8630 *to the Fourth Assessment Report of the Intergovernmental Panel on Climate*
8631 *Change*. [Parry, M.L., O.F. Canziani, J.P. Palutikof, and P.J. van der Linden,
8632 (eds.)]. Cambridge University Press, Cambridge, UK, and New York, pp. 79-131.
- 8633 **Rybczyk**, J.M. and D.R. Cahoon, 2002: Estimating the potential for submergence for two
8634 subsiding wetlands in the Mississippi River delta. *Estuaries*, **25(5)**, 985-998.
- 8635 **Salm**, R.V. and J.R. Clark, 2000: *Marine and Coastal Protected Areas*. International
8636 Union for Conservation of Nature and Natural Resources, Switzerland, 370 pp.
- 8637 **Schwab**, W.C., E.R. Thieler, J.R., Allen, D.S., Foster, B.A., Swift, and J.F. Denny, 2000:
8638 Influence of inner-continental shelf geologic framework on the evolution and
8639 behavior of the barrier island system between Fire Island inlet and Shinnecock
8640 inlet, Long Island, New York. *Journal of Coastal Research*, **16(2)**, 408-422.
- 8641 **Smith**, S.V. and R.W. Buddemeier, 1992: Global change and coral reef ecosystems.
8642 *Annual Reviews of Ecological Systematics*, **23**, 89-118.
- 8643 **Stive**, M.J.F., 2004: How important is global warming for coastal erosion? An editorial
8644 comment. *Climatic Change*, **64(1-2)**, 27-39.
- 8645 **Trenhaile**, A.S., 2001: Modeling the effect of late Quaternary interglacial sea levels on
8646 wave-cut shore platforms. *Marine Geology*, **172(3-4)**, 205-223.

- 8647 **Walkden**, M. and M. Dickson, 2008: Equilibrium erosion of soft rock shores with a
8648 shallow or absent beach under increased sea level rise. *Marine Geology*, **251(1-2)**,
8649 75-84.
- 8650 **Walkden**, M.J.A. and J.W. Hall, 2005: A predictive Mesoscale model of the erosion and
8651 profile development of soft rock shores. *Coastal Engineering*, **52(6)**, 535-563.
- 8652 **Wells**, J.T., 1995: Effects of sea level rise on coastal sedimentation and erosion. In:
8653 *Climate Change: Impact and Coastal Habitation*, [Eisma, D. (ed.)]. CRC Press,
8654 Boca Raton, FL, pp. 111-136.
- 8655 **Williams**, S.J., 2003: Coastal and marine processes. Chapter 1.1.3.2. In: *Our Fragile*
8656 *World: Challenges and Opportunities for Sustainable Development: Encyclopedia*
8657 *of Life Support Systems (EOLSS)*, [Cilek, V. (ed.)]. Developed under the
8658 auspices of the UNESCO, EOLSS Publishers, Oxford, UK, 13 pp.
8659 <<http://www.eolss.net>>
- 8660 **Williams**, S.J., K. Dodd, and K.K. Gohn, 1991: *Coasts in Crisis*. U.S. Geological Survey
8661 circular 1075. U.S. Geological Survey, [Reston, VA], 30 pp.
- 8662 **Woodworth**, P.L and D.L. Blackman, 2004: Evidence for systematic changes in extreme
8663 high waters since the mid-1970s. *Journal of Climate*, **17(6)**, 1190-1197.
- 8664

Chapter 14. A Science Strategy for Improving the Understanding of Sea-Level Rise and its Impacts on U.S. Coasts

Authors: E. Robert Thielert, USGS; K. Eric Anderson, USGS; Donald R. Cahoon, USGS; S. Jeffress Williams, USGS; Benjamin T. Gutierrez, USGS

KEY FINDINGS

- Understanding, predicting, and responding to the environmental and human effects of sea-level rise requires an integrated program of research that includes natural and social sciences.
- Monitoring of modern processes and environments could be improved by expanding the network of basic observations and observing systems, developing time series data on environmental and landscape changes, and assembling baseline data for the coastal zone.
- The historic and geologic record of coastal change should be used to improve the understanding of natural and human-influenced coastal systems, increase knowledge of sea-level rise and coastal change over the past few millennia, identify thresholds or tipping points in coastal systems, and more closely relate past changes in climate to coastal change.

- 8685 • Increases in predictive capabilities can be achieved by improving quantitative
8686 assessment methods and integrating studies of the past and present into predictive
8687 models.
- 8688 • Research on adaptation, mitigation, and avoidance measures will enable better
8689 understanding of the societal impacts of sea-level rise.
- 8690 • Decision making in the coastal zone can be supported by providing easy access to
8691 data and resources, transferring knowledge of vulnerability and risk that affect
8692 decision making, and educating the public about consequences and alternatives.
8693

8694 **14.1 INTRODUCTION**

8695 Chapter 14 identifies several major themes that present opportunities to improve the
8696 scientific understanding of future sea-level rise and its impacts on U.S. coastal regions.
8697 Advances in scientific understanding will enable the development of higher quality and
8698 more reliable information for planners and decision makers at all levels of government, as
8699 well as the public.

8700

8701 A number of recent studies have focused specifically on research needs in coastal areas.
8702 Two National Research Council (NRC) studies, *Science for Decision-making* (NRC,
8703 1999) and *A Geospatial Framework for the Coastal Zone* (NRC, 2004) contain
8704 recommendations for science activities that can be applied to sea-level rise studies. Other
8705 relevant NRC reports include *Responding to Changes in Sea Level* (NRC, 1987), *Sea*
8706 *Level Change* (NRC, 1990b) and *Abrupt Climate Change* (NRC, 2002). The Marine
8707 Board of the European Science Foundation's Impacts of Climate Change on the European

Marine and Coastal Environment (Philippart *et al.*, 2007) identified numerous research needs, many of which have application to the United States. Recent studies on global climate change by the Pew Charitable Trusts also included the coastal zone (*e.g.*, Neumann *et al.*, 2000, Panetta, 2003; Kennedy *et al.*, 2002). Other studies by the NRC (1990a, 1990b, 1990c, 2001, 2006a, 2007) and the Heinz Center (2000, 2002a, 2002b, 2006) have addressed issues relevant to the impacts of sea-level rise on the coastal zone. These reports and related publications have helped guide the development of the potential research and decision-support activities described in the following sections.

14.2 A SCIENCE STRATEGY TO ADDRESS SEA-LEVEL RISE

An integrated scientific program of sea level studies that seeks to learn from the historic and geologic past, and monitors ongoing physical and environmental changes, will improve the level of knowledge and reduce the uncertainty about potential responses of coasts, estuaries, and wetlands to sea-level rise. Outcomes of both natural and social scientific research will support decision making and adaptive management in the coastal zone. The main elements of a potential science strategy and their interrelationships are shown in Figure 14.1.




Figure 14.1 Schematic flow diagram summarizing a science strategy for improvement of scientific knowledge and decision-making capability needed to address the impacts of future sea-level rise.

Building on and complementing ongoing efforts at federal agencies and universities, a research and observation program could incorporate new technologies to address the complex scientific and societal issues highlighted in this Product. These studies could include further development of a robust monitoring program for all coastal regions, leveraging the existing network of site observations, as well as the growing array of coastal observing systems. Research should also include studies of the historic and recent

geologic past to understand how coastal systems evolved in response to past changes in sea level. The availability of higher resolution data collected over appropriate time spans, coupled with conceptual and numerical models of coastal evolution, will provide the basis for improved quantitative assessments and the development of predictive models useful for decision making. Providing ready access to interpretations from scientific research—as well as the underlying data—by means of publications, data portals, and decision-support systems will allow coastal managers to evaluate alternative strategies for mitigation, develop appropriate responses to sea-level rise, and practice adaptive management as new information becomes available.

14.2.1 Learn From the Historic and Recent Geologic Past

Studies of the recent geologic and historical record of sea-level rise and coastal and environmental change are needed to improve the state of knowledge of the key physical and biological processes involved in coastal change. As described throughout this Product, particularly in Chapters 1 through 5, significant knowledge gaps exist that inhibit useful prediction of future changes. The following research activities will help refine our knowledge of past changes and their causes.

Improve understanding of natural and human-influenced coastal systems

Significant opportunities exist to improve predictions of coastal response to sea-level rise. For example, scientists' understanding of the processes controlling rates of sediment flux in both natural and especially in human-modified coastal systems is still evolving. This is particularly true at the regional (littoral cell) scale, which is often the same scale at which

management decisions are made. As described in Chapters 3 and 6, the human impact on coastal processes at management scales is not well understood. Shoreline engineering such as bulkheads, revetments, seawalls, groins, jetties, and beach nourishment can fundamentally alter the way a coastal system behaves by changing the transport, storage, and dispersal of sediment. The same is true of development and infrastructure on mobile landforms such as the barrier islands that comprise much of the mid-Atlantic coast.

Develop better information on the effects of sea-level rise over the past 5,000 years

The foundation of modern coastal barrier island and wetland systems has evolved over the past 5,000 years as the rate of sea-level rise slowed significantly (see Chapters 1, 3, and 4). More detailed investigation of coastal sedimentary deposits is needed to understand the rates and patterns of change during this part of the recent geologic past. Advances in methods to obtain samples of the geologic record, along with improvements in analytical laboratory techniques since the early 1990s, have significantly increased the resolution of the centennial-to-millennial scale record of sea-level rise and coastal environmental change (*e.g.*, Gehrels, 1994; Gehrels *et al.*, 1996; van de Plassche *et al.*, 1998; Donnelly *et al.*, 2001; Horton *et al.*, 2006) and provide a basis for future work. Archaeological records of past sea-level change also exist in many locales, and provide additional opportunities to understand coastal change and impacts on human activity.

Understand thresholds in coastal systems that, if crossed, could lead to rapid changes to coastal and wetland systems

Several aspects of climate change studies, such as atmosphere-ocean interactions, vegetation change, sea ice extent, and glacier and ice cap responses to temperature and precipitation, involve understanding the potential for abrupt climate change or “climate surprises” (NRC, 2002; Meehl *et al.*, 2007). Coastal systems may also respond abruptly to changes in sea-level rise or other physical and biological processes (see Chapter 3, Box 3.1). Coastal regions that may respond rapidly to even modest changes in future external forcing need to be identified, as well as the important variables driving the changes. For example, limited sediment supply, and/or permanent sand removal from the barrier system, in combination with an acceleration in the rate of sea-level rise, could result in the development of an unstable state for some barrier island systems (*i.e.*, a behavioral threshold or tipping point, as described in Chapters 1 and 3). Coastal responses could result in landward migration or roll-over, or barrier segmentation. Understanding and communicating the potential for such dramatic changes in the form and rate of coastal change will be crucial for the development of adaptation, mitigation, and other strategies for addressing sea-level rise.

The future evolution of low-elevation, narrow barriers will likely depend in part on the ability of salt marshes in back-barrier lagoons and estuaries to keep pace with sea-level rise (FitzGerald *et al.*, 2004, 2008; Reed *et al.*, 2008). It has been suggested that a reduction of salt marsh in back-barrier regions could change the hydrodynamics of back-barrier systems, altering local sediment budgets and leading to a reduction in sandy materials available to sustain barrier systems (FitzGerald *et al.*, 2004, 2008).

Relate climate proxies to coastal change

Links between paleoclimate proxies (*e.g.*, atmospheric gases in ice cores, isotopic composition of marine microfossils, tree rings), sea-level rise, and coastal change should be explored. Previous periods of high sea level, such as those during the last several interglacial periods, provide tangible evidence of higher-than-present sea levels that are broadly illustrative of the potential for future shoreline changes. For example, high stands of sea level approximately 420,000 and 125,000 years ago left distinct shoreline and other coastal features on the U.S. Atlantic coastal plain (Colquhoun *et al.*, 1991; Baldwin *et al.*, 2006). While the sedimentary record of these high stands is fragmentary, opportunities exist to relate past shoreline positions with climate proxies to improve the state of knowledge of the relationships between the atmosphere, sea level, and coastal evolution. Future studies may also provide insight into how coastal systems respond to prolonged periods of high sea level and rapid sea-level fluctuations during a high stand. Examples of both exist in the geologic record and have potential application to understanding and forecasting future coastal evolution.

14.2.2 Monitor Modern Coastal Conditions

The status and trends of sea-level change, and changes in the coastal environment, are monitored through a network of observation sites, as well as through coastal and ocean observing systems. Monitoring of modern processes and environments could be improved by expanding the network of basic observations, as well as the continued development of coastal and ocean observing systems. There are numerous ongoing

efforts that could be leveraged to contribute to understanding patterns of sea-level rise over space and time and the response of coastal environments.

Expand the network of basic observations

An improvement in the coverage and quality of the U.S. network of basic sea-level observations could better inform researchers about the rate of sea-level rise in various geographic areas. Tide gauges are a primary source of information for sea-level rise data at a wide range of time scales, from minutes to centuries. These data contribute to a multitude of studies on local to global sea-level trends. Tide gauge data from the United States include some of the longest such datasets in the world and have been especially valuable for monitoring long-term trends. A denser network of high-resolution gauges would more rigorously assess regional trends and effects. The addition of tide gauges along the open ocean coast of the United States would be valuable in some regions. These data can be used in concert with satellite altimetry observations.

Tide-gauge observations also provide records of terrestrial elevation change that contributes to relative sea-level change, and can be coupled with field- or model-based measurements or estimates of land elevation changes. Existing and new gauges should be co-located with continuously operating Global Positioning System (GPS) reference stations (CORS) or surveyed periodically using GPS and other Global Navigation Satellite System technology. This will enable the coupling of the geodetic (earth-based) reference frame and the oceanographic reference frame at the land-sea interface. Long time series from CORS can provide precise local vertical land movement information in

the ellipsoidal frame (*e.g.*, Snay *et al.*, 2007; Woppelmann *et al.*, 2007). Through a combined effort of monitoring ellipsoid heights and the geoid, as well as through gravity field monitoring, changes to coastal elevations can be adequately tracked.

Develop and maintain coastal observing systems

Observing systems have become an important tool for examining environmental change. They can be place-based (*e.g.*, specific estuaries or ocean locations) or consist of regional aggregations of data and scientific resources (*e.g.*, the developing network of coastal observing systems) that cover an entire region. Oceanographic observations also need to be integrated with observations of the physical environment, as well as habitats and biological processes.

An example of place-based observing systems is the National Estuarine Research Reserve System (NERRS: <<http://www.nerrs.noaa.gov>>), a network of 27 reserves for long-term research, monitoring, education, and resource stewardship. Targeted experiments in such settings can potentially elucidate impacts of sea-level rise on the physical environment, such as shoreline change or impacts to groundwater systems, or on biological processes, such as species changes or ecosystem impacts. Important contributions can also be made by the Long Term Ecological Research sites (<<http://www.lternet.edu>>) such as the Virginia Coast Reserve in the mid-Atlantic area (part of the focus of this Product). The sites combine long-term data with current research to examine ecosystem change over time. Integration of these ecological monitoring

8870 networks with the geodetic and tide gauge networks mentioned previously would also be
8871 an important enhancement.

8872

8873 The Integrated Ocean Observing System (IOOS) (<<http://www.ocean.us>>) will bring
8874 together observing systems and data collection efforts to understand and predict changes
8875 in the marine environment. Many of these efforts can contribute to understanding
8876 changes in sea-level rise over space and time. These observing systems incorporate a
8877 wide range of data types and sources, and provide an integrated approach to ocean
8878 studies. Such an approach should enable sea-level rise-induced changes to be
8879 distinguished from the diverse processes that drive changes in the coastal and marine
8880 environment.

8881

8882 A new initiative began in 2005 with a worldwide effort to build a Global Earth
8883 Observation System of Systems (GEOSS) (<<http://www.earthobservations.org>>) over the
8884 next 10 years. GEOSS builds upon existing national, regional, and international systems
8885 to provide comprehensive, coordinated Earth observations from thousands of instruments
8886 worldwide, which have broad application to sea-level rise studies.

8887

8888 *Develop time series data to monitor environmental and landscape changes*

8889 Observations of sea level using satellite altimetry (e.g., TOPEX/Poseidon and Jason-1)
8890 have provided new and important insights into the patterns of sea-level change across
8891 space and time. Such observations have allowed scientists to examine sea-level trends
8892 and compare them to the instrumental record (Church *et al.*, 2001, 2004), as well as

predictions made by previous climate change assessments (Rahmstorf, 2007). The satellite data provide spatial coverage not available with ground-based methods such as tide gauges, and provide an efficient means for making global observations. Plans for future research could include a robust satellite observation program to ensure comprehensive coverage.

Studies of environmental and landscape change can also be expanded across larger spatial scales and longer time scales. Examples include systematic mapping of shoreline changes and coastal barriers and dunes around the United States (*e.g.*, Morton and Miller, 2005), and other national mapping efforts to document land-use and land-cover changes (*e.g.*, the NOAA Coastal Change Analysis Program:

<http://www.csc.noaa.gov/crs/lca/ccap.html>). It is also important to undertake a rigorous study of land movements beyond the point scale of tide gauges and GPS networks. For example, the application of an emerging technology—Interferometric Synthetic Aperture Radar (InSAR)—enables the development of spatially-detailed maps of land-surface displacement over broad areas (Brooks *et al.*, 2007).

Determining wetland sustainability to current and future sea-level rise requires a broader foundation of observations if they are to be applied with high confidence at regional and national scales. In addition, there is a significant knowledge gap concerning the viability or sustainability of human-impacted and restored wetlands in a time of accelerating sea-level rise. The maintenance of a network of sites that utilize surface elevation tables and soil marker horizons for measuring marsh accretion or loss will be essential in

understanding the impacts on areas of critical wetland habitat. The addition of sites to the network would aid in delineating regional variations (Cahoon *et al.*, 2006). Similar long-term studies for coastal erosion, habitat change, and water quality are also essential.

Coastal process studies require data to be collected over a long period of time in order to evaluate changes in beach and barrier profiles and track morphological changes over a time interval where there has been a significant rise in sea level. These data will also reflect the effects of storms and the sediment budget that frequently make it difficult to extract the coastal response to sea-level change. For example, routine lidar mapping updates to track morphological changes and changes in barrier island area above mean high water (*e.g.*, Morton and Sallenger, 2003), as well as dune degradation and recovery, and shore-face profile and near-shore bathymetric evolution may provide insight into how to distinguish various time and space scales of coastal change and their relationship to sea-level rise.

Time series observations can also be distributed across the landscape and need not be tied to specific observing systems or data networks. They do, however, need a means to have their data assimilated into a larger context. For example, development of new remote sensing and *in-situ* technologies and techniques would help fill critical data gaps at the land-water interface.

Assemble and update baseline data for the coastal zone

Baseline data for the coastal zone, including elevation, bathymetry, shoreline position, and geologic composition of the coast, as well as biologic and ecologic parameters such as vegetation and species distribution, and ecosystem and habitat boundaries, should be collected at high spatial resolution. As described in Chapter 2, existing 30-m (100-ft) digital elevation models are generally inadequate for meaningful mapping and analyses in the coastal zone. The use of lidar data, with much better horizontal and vertical accuracy, is essential. While some of these mapping data are being collected now, there are substantial areas around the United States that would benefit from higher quality data. More accurate bathymetric data, especially in the near-shore, is needed for site-specific analyses and to develop a complete topographic-bathymetric model of the coastal zone to be able to predict with greater confidence wave and current actions, inundation, coastal erosion, sediment transport, and storm effects.

To improve confidence in model predictions of wetland vulnerability to sea-level rise, more information is needed on: (1) maximum accretion rates (*i.e.*, thresholds) regionally and among vegetative communities; (2) wetland dynamics across larger landscape scales; (3) the interaction of feedback controls on flooding with other accretion drivers (*e.g.*, nutrient supply and soil organic matter accumulation); (4) fine-grained, cohesive sediment supplies; and (5) changing land use in the watershed (*i.e.*, altered river flows and accommodation space for landward migration of wetlands). In addition, population data on different species in near shore areas are needed to accurately judge the effects of habitat loss or transformation. More extensive and detailed areas of habitat mapping will enable preservation efforts to be focused on the most important areas.

8961

8962 **14.2.3 Predict Future Coastal Conditions**

8963 Studies of the past history of sea-level rise and coastal response, combined with extensive
8964 monitoring of present conditions, will enable more robust predictions of future sea-level
8965 rise impacts. Substantial opportunities exist to improve methods of coastal impact
8966 assessment and prediction of future changes.

8967

8968 *Develop quantitative assessment methods that identify high-priority areas needing useful*
8969 *predictions*

8970 Assessment methods are needed to identify both geographic and topical areas most in
8971 need of useful predictions of sea-level rise impacts. For example, an assessment
8972 technique for objectively assessing potential effects of sea-level rise on open coasts, the
8973 Coastal Vulnerability Index (CVI), has been employed in the United States and elsewhere
8974 (*e.g.*, Gornitz *et al.*, 1997; Shaw *et al.*, 1998; Thieler and Hammar-Klose, 1999, 2000a,
8975 2000b). Although the CVI is a fairly simplistic technique, it can provide useful insights
8976 and has found application as a coastal planning and management tool (Thieler *et al.*,
8977 2002). Such assessments have also been integrated with socioeconomic vulnerability
8978 criteria to yield a more integrative measure of community vulnerability (Boruff *et al.*,
8979 2005).

8980

8981 Projecting long-term wetland sustainability to future sea-level rise requires data on
8982 accretionary events over sufficiently long time scales that include the return periods of
8983 major storms, floods, and droughts, as well as information on the effects of wetland

elevation feedback on inundation and sedimentation processes that affect wetland vertical accretion. Numerical models can be applied to predict wetland sustainability at the local scale, but there is not sufficient data to populate these models at the regional or national scale (see Chapter 4). Given this data constraint, current numerical modeling approaches will need to improve or adapt such that they can be applied at broader spatial scales with more confidence.

Integrate studies of past and present coastal behavior into predictive models

Existing shoreline-change prediction techniques are typically based on assumptions that are either difficult to validate or too simplistic to be reliable for many real-world applications (see Appendix 2). As a result, the usefulness of these modeling approaches has been debated in the coastal science community (see Chapter 3). Newer models that include better representations of real-world settings and processes (*e.g.*, Cowell *et al.*, 1992; Stolper *et al.*, 2005; Pietrafesa *et al.*, 2007) have shown promise in predicting coastal evolution. Informing these models with improved data on past coastal changes should result in better predictions of future changes.

The process of marine transgression across the continental shelf has left an incomplete record of sea-level and environmental change. An improved understanding of the rate and timing of coastal evolution will need to draw on this incomplete record, however, in order to improve models of coastal change. Using a range of techniques, such as high-resolution seafloor and geologic framework mapping coupled with geochronologic and paleoenvironmental studies, the record of coastal evolution during the Pleistocene (1.8

million to 11,500 years ago) and the Holocene (the last 11,500 years) can be explored to identify the position and timing of former shorelines and coastal environments.

14.2.4 Improve Understanding of Societal Impacts

Research in the social sciences will be critical to understanding the potential effects on society and social systems resulting from sea-level rise.

Increase research on adaptation, mitigation, and avoidance measures

This Product describes a wide variety of potential impacts of sea-level rise, including the effects on the physical environment, biological systems, and coastal development and infrastructure. While the ability to predict future changes is currently inadequate for many decisions, adaptation, mitigation, and avoidance strategies must evolve as scientific knowledge and predictive ability increase. For example, expanded research and assessments of the economic and environmental costs of present and future actions are needed to allow a more complete analysis of the tradeoffs involved in sea-level rise decision making. In addition, opportunities to engage stakeholders such as federal agencies, states, counties, towns, non-government organizations, and private landowners in the design and implementation of sea-level rise impact and response planning should be created.

Develop multi-disciplinary groups that integrate natural and social sciences

Interdisciplinary research that combines natural and social sciences will be crucial to understanding the interplay of the physical, environmental, and societal impacts of sea-

level rise. Development of programs that facilitate such collaborations should be encouraged.

Expand institutional capacity and overcome barriers

Substantial opportunities exist to expand and improve upon the ability of institutions to respond to sea-level rise (see Chapter 10, 11, and 12). Research is needed to define the capacity needed for decision making, as well as the methods that can be best employed (e.g., command and control, economic incentive) to achieve management goals.

Overcoming the institutional barriers described in Chapter 12 is also necessary for effective response to the management challenges presented by sea-level rise.

14.2.5 Develop Coastal Decision Support Systems for Planning and Policy Making

For coastal zone managers in all levels of government, there is a pressing need for more scientific information, a reduction in the ranges of uncertainty for processes and impacts, and new methods for assessing options and alternatives for management strategies.

Geospatial information on a wide range of themes such as topography, bathymetry, land cover, population, and infrastructure, that is maintained on regular cycle will be a key component of planning for mitigation and adaptation strategies. For example, specialized themes of data such as hydric (abundantly moist) soils may be critical to understanding the potential for wetland survival in specific areas. Developing and maintaining high-resolution maps that incorporate changes in hazard type and distribution, coastal development, and societal risk will be critical. Regularly conducting vulnerability assessments and reviews will be necessary in order to adapt to changing conditions.

9053

9054 *Provide easy access to data and information resources for federal, state, local, academic,*
9055 *and public users*

9056 Understanding and acting on scientific information about sea-level rise and its impacts
9057 will depend upon common, consistent, shared databases for integrating knowledge and
9058 providing a basis for decision making. Thematic data and other value-added products
9059 should adhere to predetermined standards to make them universally accessible and
9060 transferable through internet portals. All data should be accompanied by appropriate
9061 metadata describing its method of production, extent, quality, spatial reference,
9062 limitations of use, and other characteristics (NRC, 2004).

9063

9064 An opportunity exists to develop a national effort to develop and apply data integration
9065 tools to combine terrestrial and marine data into a seamless geospatial framework. This
9066 would involve the collection of real-time tide data and the development of more
9067 sophisticated hydrodynamic models for the entire U.S. coastline, as well as the
9068 establishment of protocols and tools for merging bathymetric and topographic datasets
9069 (NRC, 2004). Modern and updated digital flood insurance rate maps (DFIRM) that
9070 incorporate future sea-level rise are needed in the coastal zone (see Chapter 9).

9071

9072 *Transfer scientific knowledge to studies of vulnerability, risk, and societal impacts*

9073 In addition to basic scientific research and environmental monitoring, a significant need
9074 exists to integrate the results of these efforts into comprehensive vulnerability and risk
9075 assessments. Tools are needed for mapping, modeling, and communicating risk to help

public agencies and communities understand and reduce their vulnerability to, and risk of, sea-level rise hazards. Social science research activities are also needed that examine societal consequences and economic impacts of sea-level rise, as well as identify institutional frameworks needed to adapt to changes in the coastal zone. For example, analyses of the economic costs of armoring shores at risk of erosion and the expected lifespan of such efforts will be required, as will studies on the durability of armored shorefronts under different sea-level rise scenarios. The physical and biological consequences of armoring shores will need to be quantified and the tradeoffs communicated. Effective planning for sea-level rise will also require integrated economic assessments on the impact to fisheries, tourism, and commerce.

Applied research in the development of coastal flooding models for the subsequent study of ecosystem response to sea-level rise is underway in coastal states such as North Carolina (Feyen *et al.*, 2006). There is also a need for focused study on the ecological impacts of sea-level rise and in how the transfer of this knowledge can be made to coastal managers for decision-making.

Develop decision support systems

County and state planners need tools to analyze vulnerabilities, explore the implications of alternative response measures, assess the costs and benefits of options, and provide decision-making support. These might take the form of guidelines, checklists, or software tools. In addition, there is a need to examine issues in a landscape or ecosystem context rather than only administrative boundaries.

9099

9100 In addition to new and maintained data, models, and research, detailed site studies are
9101 needed to assess potential impacts on a site-specific basis and provide information that
9102 allows informed decision making. Appropriate methodologies need to be developed and
9103 made available. These will have to look at a full range of possible impacts including
9104 aquifer loss by saltwater intrusion, wetland loss, coastal erosion, and infrastructure
9105 implications, as well as the impact of adaptation measures themselves. Alternative
9106 strategies of adaptive management will be required. Each locality may need a slightly
9107 different set of responses to provide a balanced policy of preserving ecosystems,
9108 protecting critical infrastructure, and adjusting to property loss or protection. Providing a
9109 science-based set of decision support tools will provide a sound basis for making these
9110 important decisions.

9111

9112 *Educate the public on consequences and alternatives*

9113 Relative to other natural hazards such as earthquakes, volcanic eruptions, and severe
9114 weather (*e.g.*, hurricanes, tornadoes) that typically occur in minutes to days, sea-level rise
9115 has a long time horizon over which effects become clear. Thus, it is often difficult to
9116 communicate the consequences of this sometimes slow process that occurs over many
9117 years. The impacts of sea-level rise, however, are already being felt across the United
9118 States (see Chapter 13). Public education will be crucial for adapting to physical,
9119 environmental, economic, and social changes resulting from sea-level rise. Research
9120 activities that result in effective means to conduct public education and outreach
9121 concerning sea-level rise consequence and alternatives should be encouraged.

9122

CHAPTER 14 REFERENCES

* Indicates non-peer reviewed literature; available upon request

Baldwin, W.E., R.A. Morton, T.R. Putney, M.P. Katuna, M.S., Harris, P.T. Gayes, N.W. Driscoll, J.F. Denny, and W.C. Schwab, 2006: Migration of the Pee Dee River system inferred from ancestral paleochannels underlying the South Carolina Grand Strand and Long Bay inner shelf. *Geological Society of America Bulletin*, **118(5/6)**, 533-549.

Boruff, B.J., C. Emrich, and S.L. Cutter, 2005: Erosion hazard vulnerability of US coastal counties. *Journal of Coastal Research*, **21(5)**, 932-942.

Brooks, B.A., M.A. Merrifield, J. Foster, C.L. Werner, F. Gomez, M. Bevis, and S. Gill, 2007: Space geodetic determination of spatial variability in relative sea level change, Los Angeles Basin. *Geophysical Research Letters*, **34**, L01611, doi:10.1029/2006GL028171.

Cahoon, D.R., P.F. Hensel, T. Spencer, D.J. Reed, K.L., McKee, and N. Saintilan, 2006: Coastal wetland vulnerability to relative sea-level rise: wetland elevation trends and process controls. In: *Wetlands and Natural Resource Management* [Verhoeven, J.T.A., B. Beltman, R. Bobbink, and D. Whigham (eds.)]. Ecological studies volume 190. Springer, Berlin and New York, pp. 271-292.

Church, J.A., J.M. Gregory, P. Huybrechts, M. Kuhn, K. Lambeck, M.T. Nhuan, D. Qin, and P.L. Woodworth, 2001: Changes in sea level. In: *Climate Change 2001: The Scientific Basis*. Contribution of Working Group 1 to the Third Assessment Report of the Intergovernmental Panel on Climate Change [Houghton, J.T., Y. Ding, D.J. Griggs, M. Noguer, P.J. van der Linden, X. Dai, K. Maskell, and C.A. Johnson (eds.)]. Cambridge University Press, Cambridge UK, and New York, pp. 639-693.

Church, J.A., N.J. White, R. Coleman, K. Lambeck, and J.X. Mitrovica, 2004: Estimates of the regional distribution of sea-level rise over the 1950-2000 period. *Journal of Climate*, **17(13)**, 2609-2625.

- 9151 **Colquhoun**, D.J., G.H. Johnson, P.C. Peebles, P.F. Huddleston, and T. Scott, 1991:
9152 Quaternary geology of the Atlantic coastal plain. In: *Quaternary Nonglacial*
9153 *Geology: Conterminous U.S.* [Morrison, R.B. (ed.)]. The Geology of North
9154 America v. K-2. Geological Society of America, Boulder, CO, pp. 629-650.
- 9155 **Cowell**, P.J., P.S. Roy, and R.A. Jones, 1992: Shoreface translation model: computer
9156 simulation of coastal-sand-body response to sea-level rise. *Mathematics and*
9157 *Computers in Simulation*, **33(5-6)**, 603-608.
- 9158 **Donnelly**, J.P., S.S. Bryant, J. Butler, J. Dowling, L. Fan, N. Hausmann, P. Newby, B.
9159 Shuman, J. Stern, K. Westover, and T. Webb III, 2001: A 700-year sedimentary
9160 record of intense hurricane landfalls in southern New England. *Geological Society*
9161 *of America Bulletin*, **113(6)**, 714-727.
- 9162 * **Feyen**, J., K. Hess, E. Spargo, A. Wong, S. White, J. Sellars, and S. Gill, 2006:
9163 Development of continuous bathymetric/topographic unstructured coastal
9164 flooding model to study sea-level rise in North Carolina. In: *Proceedings of the*
9165 *9th International Conference on Estuarine and Coastal Modeling*, Charleston,
9166 SC, October 31 – November 2, 2005. [Spaulding, M. (ed.)]. American Society of
9167 Civil Engineers, Reston, VA, pp. 338-356.
- 9168 **FitzGerald**, D.M., I.V. Buynevich, and B.A. Argow, 2004: Model of tidal inlet and
9169 barrier island dynamics in a regime of accelerated sea-level rise. *Journal of*
9170 *Coastal Research*, **Special issue 39**, 789-795.
- 9171 **FitzGerald**, D.M., M.S. Fenster, B.A. Argow, and I.V. Buynevich, 2008: Coastal
9172 impacts due to sea-level rise. *Annual Reviews of Earth and Planetary Sciences*,
9173 **36**, 601-647.
- 9174 **Gehrels**, W.R., 1994: Determining relative sea level change from salt-marsh foraminifera
9175 and plant zones on the coast of Maine, U.S.A. *Journal of Coastal Research*,
9176 **10(4)**, 990-1009.

- 9177 **Gehrels**, W.R., D.F. Belknap, and J.T. Kelley, 1996: Integrated high-precision analyses
9178 of Holocene relative sea level changes: lessons from the coast of Maine.
9179 *Geological Society of America Bulletin*, **108(9)**, 1073-1088.
- 9180 * **Gornitz**, V. T.W. Beaty, and R.C. Daniels, 1997: *A Coastal Hazards Database for the*
9181 *U.S. West Coast*. ORNL/CDIAC-81, NDP-043C. Oak Ridge National Laboratory,
9182 Oak Ridge, TN, 162 pp. <<http://cdiac.ornl.gov/ndps/ndp043c.html>>
- 9183 * **Heinz Center**, 2000: *Evaluation of Erosion Hazards*. The H. John Heinz III Center for
9184 Science, Economics, and the Environment, Washington, DC, 203 pp.
- 9185 * **Heinz Center**, 2002a: *Human Links to Coastal Disasters*. The H. John Heinz III Center
9186 for Science, Economics, and the Environment, Washington, DC, 139 pp.
- 9187 * **Heinz Center**, 2002b: *The State of the Nation's Ecosystems: Measuring the Lands,*
9188 *Waters, and Living Resources of the United States*. Cambridge University Press,
9189 New York, 270 pp.
- 9190 * **Heinz Center**, 2006: *Filling the Gap--Priority Data Needs and Key Management*
9191 *Challenges for National Reporting on Ecosystem Condition*. The H. John Heinz
9192 III Center for Science, Economics, and the Environment, Washington, DC, 110
9193 pp.
- 9194 **Horton**, B.P., R. Corbett, S.J. Culver, R.J. Edwards, and C. Hillier, 2006: Modern salt
9195 marsh diatom distributions of the Outer Banks, North Carolina, and the
9196 development of a transfer function for high resolution reconstructions of sea level.
9197 *Estuarine, Coastal, and Shelf Science*, **69(3-4)**, 381-394.
- 9198 * **Kennedy**, V.S., R.R. Twilley, J.A. Kleypas, J.H. Cowan, and S.R. Hare, 2002: *Coastal*
9199 *and Marine Ecosystems & Global Climate Change: Potential Effects on U.S.*
9200 *Resources*. Pew Center on Global Climate Change, Arlington, VA, 52 pp.
- 9201 **Meehl**, G.A., T.F. Stocker, W.D. Collins, P. Friedlingstein, A.T. Gaye, J.M. Gregory, A.
9202 Kitoh, R. Knutti, J.M. Murphy, A. Noda, S.C.B. Raper, I.G. Watterson, A.J.
9203 Weaver, and Z.-C. Zhao, 2007: Global climate projections. In: *Climate Change*

- 9204 2007: *The Physical Science Basis*. Contribution of Working Group I to the Fourth
9205 Assessment Report of the Intergovernmental Panel on Climate Change [Solomon,
9206 S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor, and H.L.
9207 Miller (eds.)]. Cambridge University Press, Cambridge, UK, and New York, pp.
9208 747-845.
- 9209 **Morton**, R.A. and T.L. Miller, 2005: *National Assessment of Shoreline Change: Part 2,*
9210 *Historical Shoreline Changes and Associated Coastal Land Loss Along the U.S.*
9211 *Southeast Atlantic Coast*. U.S. Geological Survey open-file report 2005-1401,
9212 U.S. Geological Survey, [St. Petersburg, FL], 35 pp.
9213 <<http://pubs.usgs.gov/of/2005/1401>>
- 9214 **Morton**, R.A. and A.H. Sallenger Jr., 2003: Morphological impacts of extreme storms on
9215 sandy beaches and barriers. *Journal of Coastal Research*, **19**(3), 560-573.
- 9216 * **Neumann**, J.E., G. Yohe, R. Nicholls, and M. Manion, 2000: *Sea-level Rise and Global*
9217 *Climate Change: A Review of Impacts to U.S. Coasts*. Pew Center on Global
9218 Climate Change, Arlington, VA, 38 pp. <[http://www.pewclimate.org/global-](http://www.pewclimate.org/global-warming-in-depth/all_reports/sea_level_rise)
9219 [warming-in-depth/all_reports/sea_level_rise](http://www.pewclimate.org/global-warming-in-depth/all_reports/sea_level_rise)>
- 9220 **NRC** (National Research Council), 1987: *Responding to Changes in Sea Level:*
9221 *Engineering Implications*, National Academy Press, Washington, DC, 148 pp.
- 9222 **NRC** (National Research Council), 1990a: *Managing Coastal Erosion*. National
9223 Academy Press, Washington, DC, 182 pp.
- 9224 **NRC** (National Research Council), 1990b: *Sea Level Change*. National Academy Press,
9225 Washington, DC, 256 pp.
- 9226 **NRC** (National Research Council), 1990c: *Spatial Data Needs: The Future of the*
9227 *National Mapping Program*. National Academy Press, Washington, DC, 78 pp.
- 9228 **NRC** (National Research Council), 1999: *Science for Decisionmaking: Coastal and*
9229 *Marine Geology at the U.S. Geological Survey*. National Academy Press,
9230 Washington, DC, 113 pp.

- 9231 **NRC** (National Research Council), 2001: *Sea Level Rise and Coastal Disasters:*
9232 *Summary of a Forum*. National Academies Press, Washington, DC, 24 pp.
- 9233 **NRC** (National Research Council), 2002: *Abrupt Climate Change: Inevitable Surprises*.
9234 National Academy Press, Washington, DC, 230 pp.
- 9235 **NRC** (National Research Council), 2004: *A Geospatial Framework for the Coastal Zone:*
9236 *National Needs for Coastal Mapping and Charting*. National Academies Press,
9237 Washington, DC, 149 pp.
- 9238 **NRC** (National Research Council), 2006: *Beyond Mapping: Meeting National Needs*
9239 *Through Enhanced Geographic Information Science*. National Academies Press,
9240 Washington, DC, 100 pp.
- 9241 **NRC** (National Research Council), 2007: *Mitigating Shore Erosion on Sheltered Coasts*.
9242 National Academies Press, Washington, DC, 174 pp.
- 9243 * **Panetta**, L.E., 2003: *America's Living Oceans: Charting a Course for Sea Change: A*
9244 *Report to the Nation: Recommendations for a New Ocean Policy*. Pew Oceans
9245 Commission, Arlington, VA, 145 pp.
9246 <http://www.pewtrusts.org/pdf/env_pew_oceans_final_report.pdf>
- 9247 **Philippart**, C.J.M., R. Anadón, R. Danovaro, J.W. Dippner, K.F. Drinkwater, S.J.
9248 Hawkins, G. O'Sullivan, T. Oguz, and P.C. Reid, 2007: *Impacts of Climate*
9249 *Change on the European Marine and Coastal Environment*. Marine Board
9250 position paper 9, European Science Foundation, Strasbourg, France, 84 pp.
9251 <http://www.sesame-ip.eu/doc/MB_Climate_Change_VLIZ_05031.pdf>
- 9252 **Pietrafesa**, L.J., K. Kelleher, T. Karl, M. Davidson, M. Peng, S. Bao, D. Dickey, L. Xie,
9253 H. Liu, and M. Xia, 2007: A new architecture for coastal inundation and flood
9254 warning prediction. *Marine Technology Society Journal*, **40**(4), 71-77.
- 9255 **Rahmstorf**, S., 2007: A semi-empirical approach to projecting future sea-level rise.
9256 *Science*, **315**(5810), 368-370.

- 9257 **Reed**, D.J., D. Bishara, D. Cahoon, J. Donnelly, M. Kearney, A. Kolker, L. Leonard,
9258 R.A. Orson, and J.C. Stevenson, 2008: Site-specific scenarios for wetlands
9259 accretion as sea level rises in the mid-Atlantic region. Section 2.1 in: *Background*
9260 *Documents Supporting Climate Change Science Program Synthesis and*
9261 *Assessment Product 4.1: Coastal Elevations and Sensitivity to Sea Level Rise*
9262 [Titus, J.G. and E.M. Strange (eds.)]. EPA 430R07004. U.S. Environmental
9263 Protection Agency, Washington, DC, pp. 134-174.
9264 <<http://epa.gov/climatechange/effects/coastal/background.html>>
- 9265 **Shaw**, J., R.B. Taylor, S. Solomon, H.A. Christian, and D.L. Forbes, 1998: Potential
9266 impacts of global sea-level rise on Canadian coasts. *The Canadian Geographer*,
9267 **42(4)**, 365-379.
- 9268 **Snay**, R., M. Cline, W. Dillinger, R. Foote, S. Hilla, W. Kass, J. Ray, J. Rohde, G. Sella,
9269 and T. Soler, 2007: Using global positioning system-derived crustal velocities to
9270 estimate rates of absolute sea level change from North American tide gauge
9271 records. *Journal of Geophysical Research*, **112**, B04409,
9272 doi:10.1029/2006JB004606.
- 9273 **Stolper**, D., J.H. List, and E.R. Thieler, 2005: Simulating the evolution of coastal
9274 morphology and stratigraphy with a new morphological-behavior model
9275 (GEOMBEST). *Marine Geology*, **218(1-4)**, 17-36.
- 9276 **Thieler**, E.R. and E.S. Hammar-Klose, 1999: *National Assessment of Coastal*
9277 *Vulnerability to Sea-Level Rise: Preliminary Results for the U.S. Atlantic Coast*.
9278 U.S. Geological Survey open-file report 99-593. U.S. Geological Survey, Reston,
9279 VA, 1 sheet. <<http://pubs.usgs.gov/of/of99-593/>>
- 9280 **Thieler**, E.R. and E.S. Hammar-Klose, 2000a: *National Assessment of Coastal*
9281 *Vulnerability to Sea-Level Rise: Preliminary Results for the U.S. Pacific Coast*.
9282 U.S. Geological Survey open-file report 00-178. U.S. Geological Survey, Reston,
9283 VA, 1 sheet. <<http://pubs.usgs.gov/of/2000/of00-178/>>

- 9284 **Thieler**, E.R. and E.S. Hammar-Klose, 2000b: *National Assessment of Coastal*
9285 *Vulnerability to Sea-Level Rise: Preliminary Results for the U.S. Gulf of Mexico*
9286 *Coast*. U.S. Geological Survey open-file report 00-179. U.S. Geological Survey,
9287 Reston, VA, 1 sheet. <<http://pubs.usgs.gov/of/2000/of00-179/>>
- 9288 **Thieler**, E.R., S.J. Williams, and R. Beavers, 2002: *Vulnerability of U.S. National Parks*
9289 *to Sea-Level Rise and Coastal Change*. U.S. Geological Survey fact sheet FS 095-
9290 02. [U.S. Geological Survey, Reston, VA], 2 pp. <[http://pubs.usgs.gov/fs/fs095-](http://pubs.usgs.gov/fs/fs095-02/)
9291 [02/](http://pubs.usgs.gov/fs/fs095-02/)>
- 9292 **van de Plassche**, O., K. van der Borg, and A.F.M. de Jong, 1998: Sea level-climate
9293 correlation during the past 1400 years. *Geology*, **26(4)**, 319-322.
- 9294 **Woppelmann**, G., B.M. Miguez, M.-N. Bouin, and Z. Altamimi, 2007: Geocentric sea-
9295 level trend estimates from GPS analyses at relevant tide gauges world wide,
9296 *Global and Planetary Change*, **57(3-4)**, 396-406.
- 9297